

peritoneal exudate eosinophils harvested from polymyxin B-treated guinea pigs are similar to those harvested from untreated animals [9]. Certain differences are noteworthy. Firstly, guinea pig eosinophils had a lower resting production of hydrogen peroxide, which is not due to decreased oxidative metabolism since stimulated guinea pig eosinophils had more active oxidative metabolism. Increased resting hydrogen peroxide release by human eosinophils could represent modification of peripheral blood eosinophils by circulating factors [17]. Alternatively, the site of hydrogen peroxide release in human eosinophils may be more accessible to measurement by our extracellular assay system. Stimulated guinea pig eosinophils appear to release more hydrogen peroxide than human eosinophils; this may represent more active oxidative metabolism or a more accessible site of production. The greater amount of hydrogen peroxide release following stimulation by preopsonized zymosan may reflect differences in the numbers of C3b receptors present on human and guinea pig eosinophils.

Previous studies of eosinophils have focused on quantitative differences between eosinophils and neutrophils [16-18]. Eosinophils, usually obtained from hypereosinophilic patients, were more active than neutrophils [18-20]. Preliminary studies in our laboratory note few quantitative differences between eosinophils and neutrophils. An important quantitative difference is that eosinophils do produce small amounts of hydrogen peroxide under resting conditions, whereas neutrophils have essentially no detectable hydrogen peroxide production. Eosinophils produce approximately 10-fold more hydrogen peroxide than activated peritoneal exudate macrophages, which have a relatively low level of oxidative metabolism [14,20]; this difference is more than would be expected on the basis of laboratory variation.

Differences in hydrogen peroxide release between eosinophils from normal persons and patients with hypereosinophilic syndrome were observed, consistent with our observations that metabolic heterogeneity exists among human eosinophils [12]. Unfortunately the small number of patients available for study make it difficult to draw any conclusions. They do suggest that a worthwhile question is whether elevated hydrogen peroxide production might be responsible for enhanced parasite killing [17], especially in view of our observations that circulating factors can enhance oxidative metabolism of eosinophils [21]. The demonstration that large amounts of hydrogen peroxide can be elicited after appropriate stimuli and that small amounts are continuously secreted are further support for the hypothesis that eosinophils play an important effector role in parasitic infection.

REFERENCES

1. Mahmoud AAF, Warren KS, Peters PA: A role for the eosinophil in acquired resistance to *Schistosoma mansoni* infection as determined by anti-eosinophil serum. *J Exp Med* 142:805-813, 1975
2. Olds GR, Mahmoud AAF: Role of host granulomatous response in murine schistosomiasis. *J Clin Invest* 66:1191-1199, 1980
3. Vadas M, David JR, Butterworth AE, Pisani NT, Siongok TA: A new method for the purification of human eosinophils and neutrophils and a comparison of the ability of these cells to damage *Schistosoma mansoni*. *J Immunol* 122:1228-1236, 1979
4. Kazura JW, Grove DE: Stage-specific antibody-dependent eosinophil-mediated destruction of *Trichinella spiralis*. *Nature* 274:588-589, 1978
5. Pincus SH, Butterworth AE, David JR, Robbins M, Vadas M: Antibody-dependent eosinophil mediated damage to schistosomula of *Schistosoma mansoni*: lack of requirement for oxidative metabolism. *J Immunol* 126:1794-1799, 1981
6. Babior B: Oxygen-dependent killing by phagocytes. *N Engl J Med* 298:659-668, 721-725, 1978
7. Klebanoff SJ, Jong EC, Henderson WR Jr: The eosinophil peroxidase: purification and biological properties. *The Eosinophil in Health and Disease*. Edited by AAF Mahmoud, KF Austen. New York, Grune & Stratton, 1980, pp 99-114
8. Bass D, Szejda P: Mechanisms of killing of newborn larvae of *Trichinella spiralis* by neutrophils and eosinophils. Killing by generators of hydrogen peroxide *in vitro*. *J Clin Invest* 64:1558-1564, 1979
9. Pincus SH: Comparative metabolism of guinea pig peritoneal exudate neutrophils and eosinophils. *Proc Soc Exp Biol Med* 163:482-489, 1980
10. Pincus SH: Production of eosinophil-rich guinea pig peritoneal exudates. *Blood* 52:127-135, 1977
11. Pincus SH, DiNapoli AM, Schooley WR: Superoxide production by eosinophils: activation by histamine. *J Invest Dermatol* 79:53-57, 1982
12. Pincus SH, Schooley WR, DiNapoli AM, Broder S: Metabolic heterogeneity of eosinophils from normal and hypereosinophilic patients. *Blood* 58:1175-1811, 1981
13. Root RK, Metcalfe J, Oshino N, Chance B: H_2O_2 release from human granulocytes during phagocytosis. I. Documentation, quantitation and some regulating factors. *J Clin Invest* 55:945-955, 1975
14. Nathan CF, Silverstein SC, Brukner LH, Cohn ZA: Extracellular cytotoxicity activated by macrophages and granulocytes II. Hydrogen peroxide as a mediator of cytotoxicity. *J Exp Med* 149:100-113, 1979
15. Root RK, Metcalfe JA: H_2O_2 release from human granulocytes during phagocytosis. II. Relationship to superoxide anion formation and cellular catabolism of H_2O_2 : studies with normal and cytochalasin b-treated cells. *J Clin Invest* 60:1266-1279, 1979
16. Zeiger RS, Colten HR: Histaminase release from human eosinophils. *J Immunol* 118:540-544, 1977
17. David JR, Vadas MA, Butterworth AE, deBrito PA, Carvalho E, David RA, Bila JC, Andrade ZA: Enhanced helminthotoxicity of eosinophils from patients with eosinophilia. *N Engl J Med* 303:1147-1152, 1980
18. Klebanoff SJ, Durack DT, Rosen H, Clark RA: Functional studies on human peritoneal eosinophils. *Infect Immun* 17:167-173, 1977
19. Mickenberg IP, Root RK, Wolff SM: Bactericidal and metabolic properties of human eosinophils. *Blood* 39:67-80, 1972
20. Bass DA, Grover WN, Lewis JC, Szejda P, DeChatelet LR, McCall CE: Comparison of human eosinophils from normals and patients with eosinophilia. *J Clin Invest* 66:1265-1273, 1980
21. Vadas MA, Vargos G, Nicola N, Pincus SH, Dessein A, Metcalf D, Battye FL: Eosinophil activation by colony stimulating factor in man: analysis by flow cytometry. *Blood*, In press, 1982.

The 2nd European Symposium on Psoriasis, cosponsored by the European Society for Dermatological Research and the Italian Society of Dermatology and Venereology, will be held at the Adriatico Palace Hotel, Trieste, Italy, October 7-9, 1983. Abstracts should be received by the Scientific Secretary (Dr. E. Crivellato) before May 1, 1983. Registration forms should be submitted to the Coordinator (Dr. F. Kokelj) by July 31, 1983. For further information: Dr. C. Scarpa, Clinica Dermatologica, Università degli Studi di Trieste, c/o Ospedale Maggiore, Via Stuparich 1, 34125 Trieste, Italy (040/775005).